

Universal Approximation Depth and Errors of Narrow Belief Networks with Discrete Units

Guido F. Montúfar
gfm10@psu.edu

Department of Mathematics
Pennsylvania State University
University Park, PA 16802, USA

April 1, 2013

Abstract

We generalize recent theoretical work on the minimal number of layers of narrow deep belief networks that can approximate any probability distribution on the states of their visible units arbitrarily well, from the setting of binary units (Sutskever and Hinton, 2008; Le Roux and Bengio, 2008, 2010; Montúfar and Ay, 2011) to the setting of units with finite state spaces. In particular we show that a q -ary deep belief network with $\lfloor (q^n - 1)/q(q - 1)(n - \log_q(n) - 1) \rfloor$ layers of width n is a universal approximator of distributions on $\{0, 1, \dots, q - 1\}^n$. More generally, we bound the Kullback-Leibler model approximation errors from above, depending on the network's depth and the state spaces of the units in every layer. We provide complementary results for restricted Boltzmann machines with finite state spaces.

Keywords: Deep belief network, restricted Boltzmann machine, universal approximation, representational power, Kullback-Leibler divergence, q -ary variable

1 Introduction

A *deep belief network* (Hinton et al., 2006) is a layered stochastic network with undirected bipartite interactions between the units in the top two layers, and directed bipartite interactions between the units in all other subsequent pairs of layers, directed towards the bottom layer. The top two layers form a *restricted Boltzmann machine* (Smolensky, 1986; Freund and Haussler, 1991). The entire network defines a model of probability distributions on the states of the units in the bottom layer, the *visible* layer. When the number of units in every layer has the same order of magnitude the network is called *narrow*. The *depth* refers to the number of layers. Deep belief networks have found a great number of applications in machine learning.

The representational power of neural networks in general has been studied for several decades. For instance, a well known result (Hornik et al., 1989) shows that multilayer feedforward networks with one exponentially large layer of hidden units are universal approximators. The first proof of universal approximation by deep and narrow sigmoid belief networks is due to Sutskever and Hinton (2008). They showed that a narrow sigmoid belief network with $3(2^n - 1) + 1$ layers is a universal approximator of probability distributions on $\{0, 1\}^n$. This depth bound has been improved by Le Roux and Bengio (2010) and subsequently by Montúfar and Ay (2011), as we will discuss below in more detail. These papers have focused on the universal approximation depth of networks with binary units (binary DBNs). The purpose of the present note is to generalize that work to narrow deep belief networks with finite valued units (discrete DBNs). In addition, we give bounds for the maximal Kullback-Leibler model approximation errors. We provide analogous results for restricted Boltzmann machines with discrete units (discrete RBMs).

When a DBN has L layers of widths $n_1, \dots, n_L$ and the units in the l -th layer take states $x^l = (x_1^l, \dots, x_{n_l}^l) \in \mathcal{X}^l = \mathcal{X}_1^l \times \dots \times \mathcal{X}_{n_l}^l$, then the joint distributions on the states of all units have the form

$$p(x^1, \dots, x^L) = q(x^{L-1}, x^L) \prod_{l=1}^{L-2} p_l(x^l | x^{l+1}); \quad (1)$$

where

$$q(x, y) = \exp(\mathbf{x}^\top \Theta^{L-1} \mathbf{y}) / Z(\Theta^{L-1}); \quad (2)$$

$$p_l(x|y) = \prod_{i \in [n_l]} p_{l,i}(x_i|y); \quad (3)$$

$$p_{l,i}(x_i|y) = \exp(\mathbf{x}_i^\top \Theta_i^l \mathbf{y}) / Z(\Theta_i^l \mathbf{y}). \quad (4)$$

Here $\mathbf{x} = (\mathbf{x}_1, \dots, \mathbf{x}_n)$ denotes the entry-wise one-hot representation of $x = (x_1, \dots, x_n)$. The matrix Θ^{L-1} contains the bias and interaction weights of the top two layers, the matrix Θ_i^l contains the bias and input weights of the i -th unit in layer l , and Z is in each case the partition function. The DBN probability model consists of the marginal distributions $p(x^1) = \sum_{x^2, \dots, x^L} p(x^1, \dots, x^L)$ on the states of the units in the first layer.

A model $\mathcal{M}$ of probability distributions on a finite set $\mathcal{X}$ is a *universal approximator* iff the maximum $D_{\mathcal{M}}$ of the Kullback-Leibler divergence from any distribution on $\mathcal{X}$ to the closest distribution in $\mathcal{M}$ vanishes. More formally, $D_{\mathcal{M}} := \sup_{p \in \Delta} D(p||\mathcal{M})$, where Δ is the set of all distributions on $\mathcal{X}$, and $D(p||\mathcal{M}) := \inf_{q \in \mathcal{M}} D(p||q)$, where $D(p||q) := \sum_{x \in \mathcal{X}} p(x) \log \frac{p(x)}{q(x)}$ when $\text{supp}(p) \subseteq \text{supp}(q)$ and $D(p||q) := \infty$ otherwise. We refer to $D_{\mathcal{M}}$ as the *universal* or *maximal approximation error* of $\mathcal{M}$.

For binary DBNs the following is known:

Theorem 1. *Let DBN be a binary deep belief network probability model with L layers of width $n = 2^{k-1} + k$. If $L \geq 1 + 2^S$ for some $S \in \{0, 1, \dots, 2^{k-1}\}$, then the maximum of the Kullback-Leibler divergence from any target distribution on the states $\{0, 1\}^n$ of*




Figure 1: DBN with 5 layers of width n and $q_1, \dots, q_n$ valued units in each layer.

the visible layer to its best approximation within DBN is bounded by

$$D_{\text{DBN}} \leq (2^{k-1} - S) \log(2). \quad (5)$$

In particular, this probability model is a universal approximator whenever

$$L \geq 1 + 2^{2^{k-1}}. \quad (6)$$

Note that

$$\frac{2^n}{2(n - \log_2(n))} \leq 2^{2^{k-1}} \leq \frac{2^n}{2(n - \log_2(n) - 1)}. \quad (7)$$

The universal approximation statement, eq. (6), is due to Montúfar and Ay (2011, Theorem 2). It is based on a refinement of previous work by Le Roux and Bengio (2010), who obtained the bound $L \geq 1 + \frac{2^n}{n}$ when n is a power of two. The approximation error bound, eq. (5), was discussed recently in (Montúfar and Morton, 2012, Theorem 18). It is based on the observation that DBNs lacking the universal approximation depth still may be universal approximators for subsets of visible units, which can be used to describe explicit sub-model classes of DBNs and estimate the approximation errors.

In the following we say that a unit is q valued if its state space has cardinality q , and assume without loss of generality that $q \geq 2$. Our generalization of Theorem 1 is the following. We make the simplifying assumption that all layers have the same width, and that the units in each layer have the same state spaces. See Figure 1. The results automatically hold for networks with wider hidden layers or hidden units with larger state spaces.

Theorem 2. Let DBN be a discrete deep belief network probability model with L layers of width n and $q_1, \dots, q_n$ valued units in each layer. Let m be any integer with $n \geq m \geq \prod_{j=m+2}^n q_j$ and let $q = q_1 \geq \dots \geq q_m$, $q \geq 2$. If $L \geq 2 + \frac{q^S - 1}{q - 1}$ for some $S \in \{0, 1, \dots, m\}$, then the maximum of the Kullback-Leibler divergence from any target distribution on the states of the visible layer, $\{0, 1, \dots, q_1 - 1\} \times \dots \times \{0, 1, \dots, q_n - 1\}$,

to its best approximation within DBN is bounded by

$$D_{\text{DBN}} \leq \log\left(\prod_{j \in [m-S]} q_j\right).$$

In particular, this probability model is a universal approximator whenever

$$L \geq 2 + \frac{q^m - 1}{q - 1}.$$

When all units are q -ary and the layer width is $n = q^{k-1} + k$, then the DBN model is a universal approximator whenever $L \geq 2 + \frac{q^{q^{k-1}} - 1}{q - 1}$. Note that

$$\frac{q^n - 1}{q(q-1)(n - \log_q(n))} \leq \frac{q^{q^{k-1}} - 1}{q - 1} \leq \frac{q^n - 1}{q(q-1)(n - \log_q(n) - 1)}. \quad (8)$$

Remarks

Note that, in contrast to Theorem 1, Theorem 2 does not impose conditions on the layer width n .

The set of all distributions on $\{0, 1, \dots, q-1\}^n$ has dimension $q^n - 1$. The total number of parameters of a q -ary DBN with L layers of width n is $(L-1)(n(q-1)+1)n(q-1) + n(q-1)$. Hence the model is full dimensional only if $L \geq \frac{q^n - 1}{n(q-1)(n(q-1)+2)} + 1$. This is a parameter-counting lower bound for the universal approximation depth. The sufficiency upper-bound from Theorem 2 surpasses this by roughly a factor n . Following observations show that the upper-bound could be tight nevertheless (up to small factors of n).

Probability models with hidden variables can have dimension strictly smaller than their parameter count (dimension defect). Moreover, in some cases even full dimensional models represent only very restricted classes of distributions, as has been observed for example in tree models with hidden variables. It is known that the smallest naïve Bayes model universal approximator of distributions on $\{0, 1, \dots, q-1\}^n$ has $q^{n-1}(n(q-1)+1) - 1$ parameters when q is a prime power (see Montúfar, 2013, Theorem 13). In this case the parameter counting lower-bound is $q^n / (n(q-1) + 1)$.

At the same time we believe that the approximation-error bounds are too pessimistic. However, computing tight bounds for the maximum of the Kullback-Leibler divergence remains very challenging (this is even so for simple probability models without hidden variables).

Outline of the Proof

We will prove Theorem 2 by first studying the individual parts of the network: the RBMs formed by the top two DBN layers (Section 2); the individual units with directed inputs (Section 3); the *probability sharing* that can be realized by stacks of layers (Section 4); and finally, the sets of distributions that DBNs can represent in their bottom layer (Section 5). The proof steps of Theorem 2 can be summarized as follows:

- Show that the top RBM can approximate any probability distribution with support on a set of the form $\mathcal{X}_1 \times \cdots \times \mathcal{X}_k \times \{0\} \times \cdots \times \{0\}$ arbitrarily well.
- For a $\mathcal{X}_1$ -valued unit receiving n directed inputs, show that there is a choice of parameters for which the following holds for each possible state $h_n \in \mathcal{X}_n$ of the n -th input unit: If the input vector is $(h_1, h_2, \dots, h_n)$, then the unit outputs h'_1 with probability $p^{h_n}(h'_1)$, where p^{h_n} is an arbitrary distribution on $\mathcal{X}_1$ for all $h_n \in \mathcal{X}_n$.
- Show that there is a sequence of $\frac{q^m-1}{q-1}$ stochastic maps $p(h) \mapsto p(v) = \sum_h p(v|h)p(h)$ (each of them superposing nearly qn probability multi-sharing steps), which maps the probability distributions represented by the top RBM to any probability distribution on $\mathcal{X}_1 \times \cdots \times \mathcal{X}_n$.
- Show that the DBN can represent tractable classes of probability distributions, and estimate their maximal approximation errors.

The superposition of probability sharing steps is inspired by (Le Roux and Bengio, 2010), together with the refinements of that work devised in (Montúfar and Ay, 2011). By *probability sharing* we refer to the process of transferring an arbitrary amount of probability from a state vector x' to another state vector x'' . In contrast to the binary proofs, where each layer superposes about $2n$ sharing steps, here each layer superposes about qn multi-sharing steps, whereby each multi-sharing step transfers probability from one state to $q - 1$ states (when the units are q -ary).

2 Restricted Boltzmann Machines

We denote by $\text{RBM}_{\mathcal{X}, \mathcal{Y}}$ the restricted Boltzmann machine probability model with hidden units $Y_1, \dots, Y_m$ taking states in $\mathcal{Y} = \mathcal{Y}_1 \times \cdots \times \mathcal{Y}_m$ and visible units $X_1, \dots, X_n$ taking states in $\mathcal{X} = \mathcal{X}_1 \times \cdots \times \mathcal{X}_n$. By default RBMs are defined with binary units; however, RBMs with discrete units have appeared in (Welling et al., 2005), and their representational power has been studied in (Montúfar and Morton, 2013). The following result is closely related to (Montúfar and Morton, 2013, Theorem 16):

Theorem 3. *The model $\text{RBM}_{\mathcal{X}, \mathcal{Y}}$ can approximate any mixture $p = \sum_{i=0}^m \lambda_i p_i$ arbitrarily well, where p_0 is any product distribution, and p_i is any mixture of $(|\mathcal{Y}_i| - 1)$ product distributions, for all $i \in [m]$, whereby $\text{supp}(p_i) \cap \text{supp}(p_j) = \emptyset$ for all $1 \leq i < j \leq m$.*

Here, a *product distribution* q is a distribution on $\mathcal{X}$ that factorizes as $q(x_1, \dots, x_n) = \prod_{j \in [n]} q_j(x_j)$, where q_j is a distribution on $\mathcal{X}_j$ for all $j \in [n]$. A *mixture* is a weighted sum with non-negative weights adding to one. The support of a distribution p is $\text{supp}(p) := \{x \in \mathcal{X} : p(x) > 0\}$.

Proof of Theorem 3. Let $\mathcal{E}_{\mathcal{X}}$ denote the set of strictly positive product distributions of $X_1, \dots, X_n$. Let $\mathcal{M}_{\mathcal{X}}^k$ denote the set of all mixtures of k product distributions from $\mathcal{E}_{\mathcal{X}}$. The closure $\overline{\mathcal{M}_{\mathcal{X}}^k}$ contains all mixtures of k product distributions, including those which are not strictly positive. Let $q \circ q'$ denote the renormalized entry-wise product

with $(q \circ q')(x) = q(x)q'(x) / \sum_{x' \in \mathcal{X}} q(x')q'(x')$. The model $\text{RBM}_{\mathcal{X}, \mathcal{Y}}$ can be written, up to normalization, as the set

$$\mathcal{M}_{\mathcal{X}}^{|\mathcal{Y}_1|} \circ \dots \circ \mathcal{M}_{\mathcal{X}}^{|\mathcal{Y}_m|} = \mathbb{R}_+ \mathcal{E}_{\mathcal{X}} \circ (\mathbb{1} + \mathbb{R}_+ \mathcal{M}_{\mathcal{X}}^{|\mathcal{Y}_1|-1}) \circ \dots \circ (\mathbb{1} + \mathbb{R}_+ \mathcal{M}_{\mathcal{X}}^{|\mathcal{Y}_m|-1}). \quad (9)$$

Now consider any probability distributions $p_0 \in \mathcal{E}_{\mathcal{X}}, p'_1 \in \overline{\mathcal{M}_{\mathcal{X}}^{|\mathcal{Y}_1|-1}}, \dots, p'_m \in \overline{\mathcal{M}_{\mathcal{X}}^{|\mathcal{Y}_m|-1}}$. If $\text{supp}(p'_i) \cap \text{supp}(p'_j) = \emptyset$ for all $1 \leq i < j \leq m$, then the product $(\mathbb{1} + \lambda'_1 p'_1) \circ \dots \circ (\mathbb{1} + \lambda'_m p'_m)$ is equal to $\mathbb{1} + \sum_{i \in [m]} \lambda'_i p'_i$, up to normalization. Let $\lambda'_i = \lambda_i / \lambda_0 \sum_x p'_i(x) p_0(x)$ and $p'_i(x) = p_i(x) / p_0(x)$. Then $\lambda_0 p_0 \circ (\mathbb{1} + \sum_{i \in [m]} \lambda_i p'_i) = \sum_{i=0}^m \lambda_i p_i$. Hence the mixture distribution p is contained in the closure of the RBM model. $\square$

We now discuss certain classes of probability distributions that can be approximated arbitrarily well by RBMs, depending on their hidden units.

Let $\varrho = \{A_1, \dots, A_k\}$ be a partition of $\mathcal{X}$. The *partition model* $\mathcal{P}$ with partition ϱ is the set of all probability distributions on $\mathcal{X}$ with constant value on each A_i . Geometrically, this is the simplex with vertices $\mathbb{1}_{A_i} / |A_i|$ for all $i \in [k]$. The *coarseness* of $\mathcal{P}$ is $\max_i |A_i|$. The Kullback-Leibler divergence from partition models is well understood; in particular, by (Matúš and Ay, 2003, Corollary 1):

Lemma 4. *If $\mathcal{P}$ is a partition model of coarseness c , then $D_{\mathcal{P}} = \log(c)$.*

Now, RBMs can approximate certain partition models arbitrarily well:

Lemma 5. *Let $\mathcal{P}$ be the partition model with partition blocks $\{x_1\} \times \dots \times \{x_k\} \times \mathcal{X}_{k+1} \times \dots \times \mathcal{X}_n$ for all $x_i \in \mathcal{X}_i$, for all $i \in [k]$. When $1 + \sum_{j \in [m]} (|\mathcal{Y}_j| - 1) \geq (\prod_{i \in [k]} |\mathcal{X}_i|) / \max_{j \in [k]} |\mathcal{X}_j|$, then $\mathcal{P}$ can be approximated arbitrarily well by $\text{RBM}_{\mathcal{X}, \mathcal{Y}}$.*

Proof. Any point in $\mathcal{P}$ is a mixture of the uniform distributions on the partition blocks. The partition blocks and hence the support of these mixture components are disjoint. They are product distributions, that can be written as $p_{x_1, \dots, x_k} = \prod_{i \in [k]} \delta_{x_i} \prod_{i \in [n] \setminus [k]} u_i$, where u_i denotes the uniform distribution on $\mathcal{X}_i$. For any $j \in [k]$, any mixture of the form $\sum_{x_j \in \mathcal{X}_j} \lambda_{x_j} p_{x_1, \dots, x_k}$ is also a product distribution which factorizes as

$$\left(\sum_{x_j \in \mathcal{X}_j} \lambda_{x_j} \delta_{x_j} \right) \prod_{i \in [k] \setminus \{j\}} \delta_{x_i} \prod_{i \in [n] \setminus [k]} u_i. \quad (10)$$

Hence any point in $\mathcal{P}$ is a mixture of $(\prod_{i \in [k]} |\mathcal{X}_i|) / \max_{j \in [k]} |\mathcal{X}_j|$ product distributions of the form given in eq. (10). The claim follows from Theorem 3. $\square$

Lemma 5, together with Lemma 4, implies:

Theorem 6. *If $1 + \sum_{j \in [m]} (|\mathcal{Y}_j| - 1) \geq (\prod_{i \in \Lambda} |\mathcal{X}_i|) / \max_{j \in \Lambda} |\mathcal{X}_j|$ for some $\Lambda \subseteq [n]$, then*

$$D_{\text{RBM}_{\mathcal{X}, \mathcal{Y}}} \leq \log \left(\prod_{i \in [n] \setminus \Lambda} |\mathcal{X}_j| \right).$$

In particular, the model $\text{RBM}_{\mathcal{X}, \mathcal{Y}}$ is a universal approximator whenever

$$1 + \sum_{j \in [m]} (|\mathcal{Y}_j| - 1) \geq |\mathcal{X}| / \max_{i \in [n]} |\mathcal{X}_i|.$$




Figure 2: Star graphical model of conditional distributions.

When all units are q -ary, the RBM with $(q^{n-1} - 1)/(q - 1)$ hidden units is a universal approximator of distributions on $\{0, 1, \dots, q - 1\}^n$. Theorem 6 generalizes previous results on binary RBMs (Montúfar and Ay, 2011, Theorem 1) and (Montúfar et al., 2011, Theorem 5.1), where it is shown that a binary RBM with $2^{n-1} - 1$ hidden units is a universal approximator of distributions on $\{0, 1\}^n$, and that the maximal approximation error decreases at least logarithmically in the number of hidden units. A previous result by Le Roux and Bengio (2008, Theorem 2) shows that a binary RBM with $2^n + 1$ hidden units is a universal approximator of distributions on $\{0, 1\}^n$.

3 The Internal Node of a Star

Consider an inwards directed star graph, as shown in Figure 2, with leaf variables taking states in $\mathcal{Y} = \mathcal{Y}_1 \times \dots \times \mathcal{Y}_m$, and an internal node variable taking states in $\mathcal{V}$. Denote by $\mathcal{S}_{\mathcal{Y}, \mathcal{Y}}$ the set of conditionals on $\mathcal{V}$ defined by this network, as in eq. (4):

$$p(v|y; \Theta) = \exp(\mathbf{v}^\top \Theta \mathbf{y}) / Z(\Theta \mathbf{y}), \quad \text{for all } y \in \mathcal{Y}. \quad (11)$$

The model $\mathcal{S}_{\mathcal{Y}, \mathcal{Y}}$ can represent certain stochastic maps that we will use to define a probability sharing scheme in Section 4.

Lemma 7. *Let $\mathcal{Z} = \{y_1\} \times \dots \times \{y_{k-1}\} \times \mathcal{Y}_k \times \{y_{k+1}\} \times \dots \times \{y_m\} \subseteq \mathcal{Y}$, $k \neq m$, let $\mathcal{V} = \mathcal{Y}_m$, and let $\{q^z : z \in \mathcal{Z}\}$ be any distributions on $\mathcal{V}$. Then there is a choice of the parameters Θ of $\mathcal{S}_{\mathcal{Y}, \mathcal{Y}}$ such that*

$$p(\cdot|y; \Theta) = \begin{cases} q^y, & \text{if } y \in \mathcal{Z} \\ \delta_{y_m}, & \text{otherwise} \end{cases}.$$

Proof. Let $\mathcal{Y}_j = \{0, 1, \dots, r_j - 1\}$ for all $j \in [m]$, and $r = |\mathcal{V}| = r_m$. The set of strictly positive probability distributions on $\mathcal{V}$ is an exponential family $\mathcal{E}_{\mathcal{V}} = \{p(v; \theta) = \exp(\mathbf{v}^\top \theta) / Z(\theta) : \theta \in \mathbb{R}^d\}$. Let ϑ_v be the parameter vector of a distribution which attains a unique maximum at v . Then for any fixed $\eta \in \mathbb{R}^d$ we have

$$\lim_{K \rightarrow \infty} p(x; \eta + K\vartheta_v) = \delta_v(x). \quad (12)$$

To see this, note that $p(x; K\vartheta_v) \propto p(x; \vartheta_v)^K$ and hence $\lim_{K \rightarrow \infty} p(x; K\vartheta_v) = \delta_v$, and furthermore, $p(x; \eta + K\vartheta_v) \propto p(x; \eta)p(x; K\vartheta_v)$.

Without loss of generality let $\mathcal{Z} = \mathcal{Y}_1 \times \{0\} \times \dots \times \{0\}$. For each $z \in \mathcal{Z}$ let $\theta^{z_1} \in \mathbb{R}^d$ be such that $q^z(v) = p(v; \theta^{z_1}) \in \mathcal{E}_{\mathcal{V}}$. The map Θ can be defined as follows:

$$\Theta = [\Theta_1 \mid \Theta_2 \mid \dots \mid \Theta_m]; \quad (13)$$

where Θ_j contains the columns corresponding to $\mathbf{y}_j$ in eq. (11) and

$$\begin{aligned} \Theta_1 &= [\theta^0 \mid \theta^1 \mid \dots \mid \theta^{r_1-1}] && \in \mathbb{R}^{d \times r_1}; \\ \Theta_j &= [0 \mid K_0 \vartheta_0 \mid K_0 \vartheta_0 \mid \dots \mid K_0 \vartheta_0] && \in \mathbb{R}^{d \times r_j}, \text{ for } j = 2, \dots, m-1; \\ \Theta_m &= [0 \mid K_1 \vartheta_1 \mid K_2 \vartheta_2 \mid \dots \mid K_{r-1} \vartheta_{r-1}] && \in \mathbb{R}^{d \times r}. \end{aligned} \quad (14)$$

The matrix Θ maps $\{\mathbf{y}: y \in \mathcal{Z}\}$ into the parameter vectors $\{\theta^{z_1}: z_1 \in \mathcal{Y}_1\}$ with corresponding distributions $\{q^z: z \in \mathcal{Z}\}$. When $K_0, \dots, K_{r-1} \in \mathbb{R}$ are chosen such that $\|\theta^0\|, \dots, \|\theta^{r_1-1}\| \ll K_0 \ll K_1, \dots, K_{r-1}$, then, for each $y \in \mathcal{Y} \setminus \mathcal{Z}$, the vector $\mathbf{y}$ is mapped into a parameter vector $\Theta \mathbf{y}$ with $p(\cdot | y; \Theta \mathbf{y})$ arbitrarily close to δ_{y_m} . $\square$

Remark 8. In order to prove Lemma 7 for any $\mathcal{Z}$ it is sufficient to show that (i) the vectors $\{\mathbf{y}: y \in \mathcal{Z}\}$ are affinely independent, and (ii) there is a linear map Θ mapping $\{\mathbf{y}: y \in \mathcal{Z}\}$ into the zero vector, and $\mathbf{y}$ into the relative interior of the normal cone of $Q_{\mathcal{V}} := \text{conv}\{\mathbf{v}: v \in \mathcal{V}\}$ at the vertex $v = h_m$ for all $h \in \mathcal{Y} \setminus \mathcal{Z}$.

4 Probability Sharing

A single directed layer

Consider an input layer $Y_1, \dots, Y_m$ with directed bipartite connections towards an output layer $X_1, \dots, X_n$. Denote by $\mathcal{L}_{\mathcal{X}, \mathcal{Y}}$ the model of conditional distributions defined by this network, as in eq. (3). This model defines a family of linear stochastic maps $L_{\theta}: q \mapsto \sum_{y \in \mathcal{Y}} p(x|y; \theta) q(y)$ from the simplex $\Delta(\mathcal{Y})$ of distributions on $\mathcal{Y}$ to the simplex $\Delta(\mathcal{X})$ of distributions on $\mathcal{X}$.

For any $y \in \mathcal{Y}$ and $j \in [m]$, we denote by $y[j]$ the one-dimensional cylinder set $\{y_1\} \times \dots \times \{y_{j-1}\} \times \mathcal{Y}_j \times \{y_{j+1}\} \times \dots \times \{y_m\}$. Similarly, for any $\Lambda \subseteq [m]$, we denote by $y[\Lambda]$ the set of all arrays with fixed values $\{y_i\}_{i \in [m] \setminus \Lambda}$ in the entries $[m] \setminus \Lambda$.

Applying Lemma 7 to each output unit of $\mathcal{L}_{\mathcal{X}, \mathcal{Y}}$ shows:

Theorem 9. Consider some $\{y^{(s)}\}_{s \in [k]} \subseteq \mathcal{Y}$, and let $\{j_s\}_{s \in [k]}, \{i_s\}_{s \in [k]} \subseteq [m]$. If the sets $y^{(s)}[j_s]$ are disjoint, and $\mathcal{Z}$ is a subset of $\mathcal{Y}$ containing them, then the image of $\Delta(\mathcal{Z})$ by $\mathcal{L}_{\mathcal{Y}, \mathcal{Y}}$ contains $\Delta(\mathcal{Z} \cup_{s \in [k]} y^{(s)}[\{j_s, i_s\}])$.

A stack of directed layers

In the case of binary units, sequences of probability sharing can be defined conveniently using Gray codes, as done in (Le Roux and Bengio, 2010). A Gray code is an ordered sequence of vectors, where each two subsequent vectors differ in only one entry. A binary Gray code can be viewed as a sequence of one-dimensional cylinder sets. In the

00	01	02	03	$(0, \mathcal{Y}_2)$
10	11	12	13	$(1, \mathcal{Y}_2)$
20	21	22	23	

$(\mathcal{Y}_1, 0)$

Figure 3: Three multi-sharing steps on $\{0, 1, 2\} \times \{0, 1, 2, 3\}$.

non-binary case, this correspondence is no longer present. Motivated by Theorem 9, we will use one-dimensional cylinder sets to define sequences of multi-sharing steps, as shown in Figure 3.

Let $q_i = |\mathcal{Y}_i|$ denote the cardinality of $\mathcal{Y}_i$ for $i \in [n]$, and let $m \leq n$. The set $\mathcal{Z} = \{0\} \times \cdots \times \{0\} \times \mathcal{Y}_{m+1} \times \cdots \times \mathcal{Y}_n \subseteq \mathcal{Y}$ can be written as the disjoint union of $k = \prod_{i=m+2}^n |\mathcal{Y}_i|$ one-dimensional cylinder sets, as $\mathcal{Z} = \cup_{s=1}^k y^{(s)}[m+1]$, where $y^{(s)} = (0, \dots, 0 \mid 0, y_{m+2}^{(s)}, \dots, y_n^{(s)})$ and $\{(y_{m+2}^{(s)}, \dots, y_n^{(s)})\}_{s=1}^k = \mathcal{Y}_{m+2} \times \cdots \times \mathcal{Y}_n$.

In the following, each set $y^{(s)}[m+1]$ will be the starting point of a sequence of sharing steps. By Theorem 9, a directed DBN layer can map the simplex of distributions $\Delta(y^{(1)}[m+1] \cup \cdots \cup y^{(k)}[m+1])$ surjectively onto the simplex of distributions $\Delta(y^{(1)}[m+1, 1] \cup \cdots \cup y^{(k)}[m+1, k])$. The latter can be mapped by a further DBN layer onto a larger simplex and so forth. Starting with $y^{(1)}[m+1]$, consider the sequence

$$\begin{aligned}
& (0, 0, \dots, 0 \mid 0, y_{m+2}^{(1)}, \dots, y_n^{(1)})[m+1, 1] \\
& (0, 0, \dots, 0 \mid 0, y_{m+2}^{(1)}, \dots, y_n^{(1)})[m+1, 2] \\
& (1, 0, \dots, 0 \mid 0, y_{m+2}^{(1)}, \dots, y_n^{(1)})[m+1, 2]
\end{aligned} \tag{15}$$

continued as shown in Table 1. We denote this sequence by G^1 , and its l -th row (a cylinder set) by $G^1(l)$. The union $\cup_{l \in [K]} G^1(l)$ of the first K rows, with $K = 1 + q_1 + q_1 q_2 + \cdots + \prod_{j=1}^{m-1} q_j$, is equal to $\mathcal{Y}_1 \times \cdots \times \mathcal{Y}_m \times \mathcal{Y}_{m+1} \times \{y_{m+2}^{(1)}\} \times \cdots \times \{y_n^{(1)}\}$.

We define k sequences $G^1, \dots, G^k$ as follows: The first m coordinates of G^s are equal to a permutation of the first m coordinates of G^1 , shifting each column cyclically s positions to the right. The last $n-m$ coordinates of G^s are equal to $(\mathcal{Y}_{m+1}, y_{m+2}^{(s)}, \dots, y_n^{(s)})$.

We abbreviate $\{s+t\} := (s+t-1) \bmod (m) + 1$. Within the first m columns, the free coordinate of the l -th row of G^s is $s + \kappa$, where κ is the least integer with $l \leq \sum_{i=0}^{\kappa} \prod_{j=s}^{\{s+i-1\}} q_j$. Here the empty product is defined as 1. Let $q = \max_{j \in \{1, \dots, m\}} q_j$. We can modify each sequence G^s , by repeating rows if necessary, such that the free coordinate of the l -th row of the resulting sequence $\tilde{G}^s$ is $s + \kappa$, where κ is the least integer with $l \leq \sum_{t=0}^{\kappa} q^t$. This κ does not depend on s .

The sequences $\tilde{G}^s$ for $s \in \{1, \dots, k\}$, are all different from each other in the last $n-m$ coordinates and have a different ‘sharing’ free-coordinate in each row. The union of cylinder sets in all rows of these sequences is equal to $\mathcal{Y}_1 \times \cdots \times \mathcal{Y}_n$.

$\mathcal{Y}_1$	0	0	0	0	
0	$\mathcal{Y}_2$	0	0	0	
1	$\mathcal{Y}_2$	0	0	0	
$\vdots$	$\vdots$				
$q_1 - 1$	$\mathcal{Y}_2$	0	0	0	
0	0	$\mathcal{Y}_3$	0	0	
0	1	$\mathcal{Y}_3$	0	0	
$\vdots$	$\vdots$	$\vdots$			
0	$q_2 - 1$	$\mathcal{Y}_3$	0	0	
1	0	$\mathcal{Y}_3$	0	0	
1	1	$\mathcal{Y}_3$	0	0	
$\vdots$	$\vdots$	$\vdots$			
1	$q_2 - 1$	$\mathcal{Y}_3$	0	0	
$\vdots$	$\vdots$				
$q_1 - 1$	0	$\mathcal{Y}_3$	0	0	
$q_1 - 1$	1	$\mathcal{Y}_3$	0	0	
$\vdots$	$\vdots$	$\vdots$			
$q_1 - 1$	$q_2 - 1$	$\mathcal{Y}_3$	0	0	
0	0	0	$\mathcal{Y}_4$	0	
0	0	1	$\mathcal{Y}_4$	0	
$\vdots$	$\vdots$	$\vdots$	$\vdots$		
0	0	$q_3 - 1$	$\mathcal{Y}_4$	0	
$\vdots$	$\vdots$				
$q_1 - 1$	$q_2 - 1$	$\cdots$	$q_{m-2} - 1$	0	$\mathcal{Y}_m$
$q_1 - 1$	$q_2 - 1$	$\cdots$	$q_{m-2} - 1$	1	$\mathcal{Y}_m$
$\vdots$	$\vdots$		$\vdots$	$\vdots$	$\vdots$
$q_1 - 1$	$q_2 - 1$	$\cdots$	$q_{m-2} - 1$	$q_{m-1} - 1$	$\mathcal{Y}_m$

Table 1: Sequence of one-dimensional cylinder sets.

5 Deep Belief Networks

Proposition 10. Consider a DBN with $L \geq 2$ layers of width n , each layer containing units with state spaces of cardinalities $q_1, \dots, q_n$. Let m be any integer with $n \geq m \geq \prod_{j=m+2}^n q_j =: k$. This model can approximate a distribution p arbitrarily well whenever the support of p is contained in $\cup_{s \in [k]} \cup_{l \in [L-2]} \tilde{G}^s(l)$.

Proof. Note that $\prod_{j=m+2}^n q_j \leq n \leq 1 + \sum_{j \in [n]} (q_j - 1)$. By Theorem 3 the top RBM can approximate the probability simplex on $\{0\} \times \dots \times \{0\} \times \mathcal{X}_{m+1} \times \dots \times \mathcal{X}_n$ arbitrarily well. By Theorem 9, this simplex can be mapped iteratively into larger simplices, according to the sequences $\tilde{G}^s$ from Section 4. $\square$

Theorem 11. Consider a DBN with L layers of width n , each layer containing units with state spaces of cardinalities $q_1, \dots, q_n$. Let m be any integer with $n \geq m \geq \prod_{j=m+2}^n q_j$ and $q := q_1 \geq \dots \geq q_m$. If $L \geq 2 + 1 + q + \dots + q^{k-1} = 2 + \frac{q^k - 1}{q - 1}$, then this model can approximate a partition model $\mathcal{P}$ of coarseness $\prod_{j=1}^{m-k} q_j$ arbitrarily well.

Proof. We use the abbreviation $\{s + t\} := (s + t - 1) \bmod (m) + 1$. The top RBM can approximate a partition model $\mathcal{P}$ (on a subset of $\mathcal{Y}$) arbitrarily well, whose partition blocks are the cylinder sets with fixed coordinate values $y_s = 0, y_{\{s+1\}} = 0, y_{\{s+k\}} = 0, y_{m+1}, y_{m+2}^{(s)}, \dots, y_n^{(s)}$ for all $y_{m+1} \in \mathcal{Y}_{m+1}$, for all $s \in [k]$. When $L \geq 2 + 1 + q + q^2 + \dots + q^r$, after $L - 2$ probability sharing steps starting from $\mathcal{P}$, the DBN can approximate the partition model arbitrarily well, whose partition blocks are the cylinder sets with fixed coordinate values

$$y_s, y_{\{s+1\}} \dots, y_{\{s+r\}}, y_{m+1}, y_{m+2}^{(s)}, \dots, y_n^{(s)}; \quad (16)$$

for all possible choices of $y_s, y_{\{s+1\}}, \dots, y_{\{s+r\}}, y_{m+1}$, for all $s \in [k]$. The maximal cardinality of such a block is $q_1 \dots q_{m-r-1}$, and the union of all blocks equals $\mathcal{Y}$. $\square$

Proof of Theorem 2. The claim follows using Lemma 4 on the partition models described in Theorem 11. $\square$

Acknowledgment

The author is supported in part by DARPA grant FA8650-11-1-7145.

References

- Y. Freund and D. Haussler. Unsupervised learning of distributions of binary vectors using 2-layer networks. In *Advances in Neural Information Processing Systems 4*, pages 912–919, 1991.
- G. E. Hinton, S. Osindero, and Y. Teh. A fast learning algorithm for deep belief nets. *Neural Computation*, 18:1527–1554, 2006.

- K. Hornik, M. B. Stinchcombe, and H. White. Multilayer feedforward networks are universal approximators. *Neural Networks*, 2(5):359–366, 1989.
- N. Le Roux and Y. Bengio. Representational power of restricted Boltzmann machines and deep belief networks. *Neural Computation*, 20(6):1631–1649, 2008.
- N. Le Roux and Y. Bengio. Deep belief networks are compact universal approximators. *Neural Computation*, 22:2192–2207, 2010.
- F. Matúš and N. Ay. On maximization of the information divergence from an exponential family. In *Proceedings of the WUPES’03*, pages 199–204, 2003.
- G. Montúfar. Mixture decompositions of exponential families using a decomposition of their sample spaces. *Kybernetika*, 49(1):23–39, 2013.
- G. Montúfar and N. Ay. Refinements of universal approximation results for deep belief networks and restricted Boltzmann machines. *Neural Computation*, 23(5):1306–1319, 2011.
- G. Montúfar and J. Morton. Kernels and submodels of deep belief networks. 2012. <http://arxiv.org/abs/1211.0932>.
- G. Montúfar and J. Morton. Discrete restricted Boltzmann machines. 2013. <http://arxiv.org/abs/1301.3529>.
- G. Montúfar, J. Rauh, and N. Ay. Expressive power and approximation errors of restricted Boltzmann machines. In *Advances in Neural Information Processing Systems 24*, pages 415–423, 2011.
- P. Smolensky. Information processing in dynamical systems: foundations of harmony theory. In *Parallel distributed processing: explorations in the microstructure of cognition*, vol. 1, pages 194–281. MIT Press, 1986.
- I. Sutskever and G. E. Hinton. Deep narrow sigmoid belief networks are universal approximators. *Neural Computation*, 20:2629–2636, 2008.
- M. Welling, M. Rosen-Zvi, and G. E. Hinton. Exponential family harmoniums with an application to information retrieval. In *Advances in Neural Information Processing Systems 17*, pages 1481–1488, 2005.